

**INBRED CORN LINE G3001****FIELD OF THE INVENTION**

5           This invention is in the field of corn breeding, specifically relating to an inbred corn line designated G3001. This invention also is in the field of hybrid maize production employing the present inbred.

**BACKGROUND OF THE INVENTION**

10           The original maize plant was indigenous to the Western Hemisphere. The plants were weed-like and only through the efforts of early breeders were cultivated crop species developed. The crop cultivated by early breeders, like the crop today, could be wind pollinated. The physical traits of maize are such that  
15           wind pollination results in self-pollination or cross-pollination between plants. Each plant has a separate male and female flower that contributes to pollination, the tassel and ear, respectively. Natural pollination occurs when wind transfers pollen from tassel to the silks on the corn ears. This type of pollination has contributed to the wide variation of maize varieties present in the Western Hemisphere.

20           The development of a planned breeding program for maize only occurred in the last century. A large part of the development of the maize product into a profitable agricultural crop was due to the work done by land grant colleges. Originally, maize was an open pollinated variety having heterogeneous genotypes.  
25           The maize farmer selected uniform ears from the yield of these genotypes and preserved them for planting the next season. The result was a field of maize plants that were segregating for a variety of traits. This type of maize selection led to, at most, incremental increases in seed yield.

30           Large increases in seed yield were due to the work done by land grant

colleges that resulted in the development of numerous hybrid corn varieties in planned breeding programs. Hybrids were developed by selecting corn lines and selfing these lines for several generations to develop homozygous pure inbred lines. One selected inbred line was crossed with another selected inbred line to produce hybrid progeny (F1). Although hybrids, due to heterosis, are robust and vigorous plants. Inbreds on the other hand are mostly homozygous. This homozygosity renders the inbred lines less vigorous. Inbred seed can be difficult to produce since the inbreeding process in corn lines decreases the vigor. However, when two inbred lines are crossed, the hybrid plant evidences greatly increased vigor and seed yield compared to open pollinated, segregating maize plants. An important consequence of the homozygosity and the homogeneity of the inbred maize lines is that all hybrid seed produced from any cross of two such elite lines will be the same hybrid seed and make the same hybrid plant. Thus the use of inbreds makes hybrid seed which can be reproduced readily. The hybrid plant in contrast does not produce hybrid seed that is readily reproducible. The seed on a hybrid plant is segregating for traits.

The ultimate objective of the commercial maize seed companies is to produce high yielding, agronomically sound plants that perform well in certain regions or areas of the Corn Belt. To produce these types of hybrids, the companies must develop inbreds, which carry needed traits into the hybrid combination. Hybrids are not often uniformly adapted for the entire Corn Belt, but most often are specifically adapted for regions of the Corn Belt. Northern regions of the Corn Belt require shorter season hybrids than do southern regions of the Corn Belt. Hybrids that grow well in Colorado and Nebraska soils may not flourish in richer Illinois and Iowa soil. Thus, a variety of major agronomic traits are important in hybrid combination for the various Corn Belt regions, and have an impact on hybrid performance.

Inbred line development and hybrid testing have been emphasized in the past half-century in commercial maize production as a means to increase hybrid performance. Inbred development is usually done by pedigree selection. Pedigree selection can be selection in an  $F_2$  population produced from a planned cross of two genotypes (often elite inbred lines), or selection of progeny of synthetic varieties, open pollinated, composite, or backcrossed populations. This type of selection is effective for highly inheritable traits, but other traits, for example, yield requires replicated test crosses at a variety of stages for accurate selection.

Maize breeders select for a variety of traits in inbreds that impact hybrid performance along with selecting for acceptable parental traits. Such traits include: yield potential in hybrid combination; dry down; maturity; grain moisture at harvest; greensnap; resistance to root lodging; resistance to stalk lodging; grain quality; disease and insect resistance; ear and plant height. Additionally, Hybrid performance will differ in different soil types such as low levels of organic matter, clay, sand, black, high pH, low pH; or in different environments such as wet environments, drought environments, and no tillage conditions. These traits appear to be governed by a complex genetic system that makes selection and breeding of an inbred line extremely difficult. Even if an inbred in hybrid combination has excellent yield (a desired characteristic), it may not be useful because it fails to have acceptable parental traits such as seed yield, seed size, pollen production, good silks, plant height, etc.

To illustrate the difficulty of breeding and developing inbred lines, the following example is given. Two inbreds compared for similarity of 29 traits differed significantly for 18 traits between the two lines. If 18 simply inherited single gene traits were polymorphic with gene frequencies of 0.5 in the parental lines, and assuming independent segregation (as would essentially be the case if

each trait resided on a different chromosome arm), then the specific combination of these traits as embodied in an inbred would only be expected to become fixed at a rate of one in 262,144 possible homozygous genetic combinations. Selection of the specific inbred combination is also influenced by the specific selection environment on many of these 18 traits which makes the probability of obtaining this one inbred even more remote. In addition, most traits in the corn genome are regrettably not single dominant genes but are multi-genetic with additive gene action not dominant gene action. Thus, the general procedure of producing a non segregating  $F_1$  generation and self pollinating to produce a  $F_2$  generation that segregates for traits and selecting progeny with the visual traits desired does not easily lead to a useful inbred. Great care and breeder expertise must be used in selection of breeding material to continue to increase yield and the agronomics of inbreds and resultant commercial hybrids.

Certain regions of the Corn Belt have specific difficulties that other regions may not have. Thus the hybrids developed from the inbreds have to have traits that overcome or at least minimize these regional growing problems. Examples of these problems include in the eastern corn belt Gray Leaf Spot, in the north cool temperatures during seedling emergence, in the Nebraska region CLN (corn Lethal necrosis and in the west soil that has excessively high pH levels. The industry often targets inbreds that address these issues specifically forming niche products. However, the aim of most large seed producers is to provide a number of traits to each inbred so that the corresponding hybrid can be useful in a broader regions of the Corn Belt. The new biotechnology techniques such as Microsatellites, RFLPs, RAPDs and the like have provided breeders with additional tools to accomplish these goals.

## SUMMARY OF THE INVENTION

5 The present invention relates to an inbred corn line G3001. Specifically, this invention relates to plants and seeds of this line. Additionally, this relates to a method of producing from this inbred, hybrid seed corn and hybrid plants with seeds from such hybrid seed. More particularly, this invention relates to the unique combination of traits that combine in corn line G3001.

10 Generally then, broadly the present invention includes an inbred corn seed designated G3001. This seed produces a corn plant.

15 The invention also includes the tissue culture of regenerable cells of G3001 wherein the cells of the tissue culture regenerates plants capable of expressing the genotype of G3001. The tissue culture is selected from the group consisting of leaves, pollen, embryos, roots, root tips, guard cells, ovule, seeds, anthers, silk, flowers, kernels, ears, cobs, husks and stalks, cells and protoplasts thereof. The corn plant regenerated from G3001 or any part thereof is included in the present invention. The present invention includes regenerated corn plants that are capable of expressing G3001's genotype, phenotype or mutants or variants thereof.

20 The invention extends to hybrid seed produced by planting, in pollinating proximity which includes using preserved maize pollen as explained in U.S. Patent 5,596,838 to Greaves , seeds of corn inbred lines G3001 and another inbred line if preserved pollen is not used; cultivating corn plants resulting from said planting; preventing pollen production by the plants of one of the inbred lines if two are employed; allowing cross pollination to occur between said inbred lines; and harvesting seeds produced on plants of the selected inbred. The hybrid seed produced by hybrid combination of plants of inbred corn seed designated G3001

and plants of another inbred line are apart of the present invention. This inventions scope covers hybrid plants and the plant parts including the grain and pollen grown from this hybrid seed.

5           The invention further includes a method of hybrid F1 production. A first generation (F1) hybrid corn plant produced by the process of planting seeds of corn inbred line G3001; cultivating corn plants resulting from said planting; permitting pollen from another inbred line to cross pollinate inbred line G3001; harvesting seeds produced on plants of the inbred; and growing a harvested seed  
10           are part of the method of this invention.

          Likewise included is a first generation (F1) hybrid corn plant produced by the process of planting seeds of corn inbred line G3001; cultivating corn plants resulting from said planting; permitting pollen from inbred line G3001 to cross  
15           pollinate another inbred line; harvesting seeds produced on plants of the inbred; and growing a plant from such a harvested seed.

          The inbred corn line G3001 and at least one transgenic gene adapted to give G3001 additional and/or altered phenotypic traits are within the scope of the invention. Such transgenes are usually associated with regulatory elements  
20           (promoters, enhancers, terminators and the like). Presently, transgenes provide the invention with traits such as insect resistance, herbicide resistance, disease resistance increased or decreased starch or sugars or oils, increased or decreased life cycle or other altered trait.

25           The present invention includes inbred corn line G3001 and at least one transgenic gene adapted to give G3001 modified starch traits. Furthermore this invention includes the inbred corn line G3001 and at least one mutant gene adapted to give modified starch, acid or oil traits. The present invention includes

the inbred corn line G3001 and at least one transgenic gene selected from the group consisting of: bacillus thuringiensis, the bar or pat gene encoding Phosphinothricin acetyl Transferase, Gdha gene, EPSP synthase gene, low phytic acid producing gene, and zein. The inbred corn line G3001 and at least one transgenic gene useful as a selectable marker or a screenable marker are covered by the present invention.

A tissue culture of the regenerable cells of hybrid plants produced with use of G3001 genetic material is covered by this invention. A tissue culture of the regenerable cells of the corn plant produced by the method described above are also included.

## DEFINITIONS

In the description and examples, which follow, a number of terms are used. In order to provide a clear and consistent understanding of the specifications and claims, including the scope to be given such terms, the following definitions are provided.

### **BL MOIST**

The moisture percentage of the grain at black layer, i.e., when 50% of the plants per plot have reached physiological maturity.

### **COLD GERM**

Cold Germ is a measurement of seed germination under cold soil conditions. Data is reported as percent of seed germinating.

### **ECB**

European corn borer is a maize eating insect. ECBI is the first brood generation of

European corn borers. ECBII is the second generation of European corn borers.

## **EMERGE (EMG)**

The number of emerged plants per plot (planted at the same seedling rate) collected when plants have two fully developed leaves.

## **GI**

This is a selection index that provides a single quantitative measure of the worth of a hybrid based on four traits. FI is a very similar index which weights yield less than GI. In GI yield is the primary trait contributing to index values. The GI value is calculated by combining stalk lodging, root lodging, yield and dropped ears according to the attached mathematical formula:

$$GI = 100 + 0.5 (YLD) - 0.9 (\%STALK LODGE) - 0.9 (\%ROOT LODGE) - 2.7 (\%DROPPED EAR)$$

## **GLS**

Gray Leaf Spot (*Cercospora Zeae*) disease rating. This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.\*

## **GW**

Goss' Wilt (*Corynebacterium nebraskense*). This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.\*

## HEATP10

The number of Growing Degree Units (GDU's) or heat units required for an inbred line or hybrid to have approximately 10 percent of the plants shedding pollen. This trait is measured from the time of planting. Growing Degree Units are calculated by the Barger Method where the GDU's for a 24 hour period are:

$$\text{GDU} = \frac{(\text{Max Temp } (^{\circ}\text{F}) + \text{Min Temp } (^{\circ}\text{F})) - 50}{2}$$

The highest maximum temperature used is 86°F and the lowest minimum temperature used is 50°F. For each inbred or hybrid it takes a certain number of GDU's to reach various stages of plant development.

## HEATBL

The number of GDU's after planting when approximately 50 percent of the inbred or hybrid plants in a plot have grain that has reached physiological maturity (black layer).

## HEATPEEK

The number of GDU's after planting of an inbred when approximately 50 percent of the plants show visible tassel extension.

## HEATP50 or HTP50

The number of GDU's required for an inbred or hybrid to have approximately 50 percent of the plants shedding pollen. Growing Degree Units are calculated by the

Barger Method as shown in the HEATP10 definition.

### **HEATP90**

5        The number of GDU's accumulated from planting when the last 100 percent of plants in an inbred or hybrid are still shedding enough viable pollen for pollination to occur. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

### 10        **HEATS10**

The number of GDU's required for an inbred or hybrid to have approximately 10 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

### 15        **HEATS50 or HTS50**

The number of GDU's required for an inbred or hybrid to have approximately 50 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

### 20        **HEATS90**

The number of GDU's required for an inbred or hybrid to have approximately 90 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

### 25        **MDMV<sub>A</sub>**

Maize Dwarf Mosaic Virus strain A. The corn is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.\*

#### **MDMV<sub>B</sub>**

Maize Dwarf Mosaic Virus strain B. This is rated on a 1-9 scale with a "1" being very susceptible and a "9" being very resistant.\*

#### **MOISTURE**

The average percentage grain moisture of an inbred or hybrid at harvest time.

#### **NLB**

Northern Leaf Blight (*Exserohilum turcicum*) disease rating. This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.\*

#### **PCT TILLER**

The total number of tillers per plot divided by the total number of plants per plot.

#### **PLANT**

This term includes plant cells, plant protoplasts, plant cell tissue cultures from which corn plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk and the like., and this term also includes any transgenic DNA or (RNA) or portion thereof that have been introduced into the plant by whatever method.

## **PLANT HEIGHT (PLTHT) (PHT)**

The distance in centimeters from ground level to the base of the tassel peduncle.

5

## **PLANT INTEGRITY (PLTINT) or (INT)**

The level of plant integrity on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are low plant integrity, 3-5.9 ratings are intermediate plant integrity, and 6-9 ratings are strongly evidencing plant integrity.

10

## **POPULATION (POP)**

The plant population.

## **RM**

Predicted relative maturity based on the moisture percentage of the grain at harvest. This rating is based on known set of checks and utilizes standard linear regression analyses and is referred to as the Minnesota Relative Maturity Rating System.

20

## **SHED**

The volume of pollen shed by the male flower rated on a 1-9 scale where a "1" is a very light pollen shedder, a "4.5" is a moderate shedder, and a "9" is a very heavy shedder.

25

## **SLB**

Southern Leaf Blight (*Bipolaris maydis*) disease rating. This is rated on a 1-9 scale

with a "1" being very susceptible, and a "9" being very resistant.\*

### **STAYGREEN (SGN)**

5 The level of staygreen of the plant on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are low staygreen, 3-5.9 ratings are intermediate staygreen, and 6-9 ratings are strongly evidencing staygreen.

### **TWT**

10 The measure of the weight of grain in pounds for a one bushel volume adjusted for percent grain moisture.

### **VIGOR (VIG)**

15 Visual rating of 1 to 9 made 2-3 weeks post-emergence where a "1" indicates very poor early plant development, and a "9" indicates superior plant development.

### **WARM GERM**

20 A measurement of seed germination under ideal (warm, moist) conditions. Data is reported as percent of seeds germinating.

### **YIELD (YLD)**

25 Actual yield of grain at harvest adjusted to 15.5% moisture. Measurements are reported in bushels per acre.

## **% DROPPED EARS (DE)**

The number of plants per plot, which dropped their primary ear, divided by the total number of plants per plot.

5

## **% LRG FLAT**

Percentage by weight of shelled corn that passes through a 26/64 inch round screen and a 14/64 inch slot screen, but does not pass through a screen with 20.5/64 inch round openings.

10

## **% LRG ROUND**

Percentage by weight of shelled corn that passes through a 26/64 inch round screen, but does not pass through a 14/64 inch slot screen or a screen with 20.5/64 inch round openings.

15

## **% MED FLAT**

Percentage by weight of shelled corn that passes through a 20.5/64 inch round screen and a 13/64 inch slotted screen, but does not pass through a screen with 17/64 inch round openings.

20

## **% MED ROUND**

Percentage by weight of shelled corn that passes through a 20.5/64 inch round screen, but does not pass through a 13/64 inch slot screen or a screen with 17/64 inch round openings.

25

### **% SML FLAT**

Percentage by weight of shelled corn that passes through a 17/64 inch round screen and a 12/64 inch slotted screen, but does not pass through a screen with 15/64 inch round openings.

### **% SML ROUND**

Percentage by weight of shelled corn that passes through a 17/64 inch round screen, but does not pass through a 12/64 inch slotted screen or a screen with 15/64 inch round openings.

### **% ROOT LODGE (RL)**

Percentage of plants per plot leaning more than 30 degrees from vertical divided by total plants per plot.

### **% STALK LODGE (SL)**

Percentage of plants per plot with the stalk broken below the primary ear node divided by the total plants per plot.

\*Resistant - on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are susceptible, 3-5.9 ratings are intermediate, and 6-9 ratings are resistant.

## DETAILED DESCRIPTION OF THE INVENTION

G3001 can be used as a female line. This inbred is useful as a female because it has nice seed producing traits. Good yields with a large percentage of the seeds being medium sized flat or rounds. In hybrid combination this line yields nice 108 day hybrids. The inbred carries a slightly better than average plant integrity into late season.

The inbred has shown uniformity and stability within the limits of environmental influence for all the traits as described in the Variety Description Information (Table 1) that follows.

The inbred has been self-pollinated for a sufficient number of generations to give inbred uniformity. During plant selection in each generation, the uniformity of plant type was selected to ensure homozygosity and phenotypic stability. The line has been increased in isolated farmland environments with data on uniformity and agronomic traits being observed to assure uniformity and stability. No variant traits have been observed or are expected in G3001.

The best method of producing the invention, G3001 which is substantially homozygous, is by planting the seed of G3001 which is substantially homozygous and self-pollinating or sib pollinating the resultant plant in an isolated environment, and harvesting the resultant seed.

TABLE 1

G3001

VARIETY DESCRIPTION INFORMATION

- #1 Type: Dent
- #2 This inbred has RM of 107.

### #3 Plant Traits

5

Plant Height 76 in.  
 Ear Height 29 in.  
 Tillers (Rating) 5  
 Leaf Color MEDIUM GREEN  
 Brace Root Color RED  
 Silk Color RED  
 Shoots at Flowering BALD

10

### #4 Tassel Traits

Glume Color GREEN/PURPLE  
 Glume Ring Color RED  
 Anther Color YELLOW

15

### #5 Ear and Kernel Traits

Cob Color RED  
 Kernel Crown Color LIGHT YELLOW  
 Kernel Body Color DARK YELLOW

20

25

## # 6 DISEASE RESISTANCE IN INBRED G3001

Eye Spot = 5.1

Gray Leaf Spot = 4.00

Gross wilt = 2.5

Northern leaf blight = 6.0

## #7 INSECT RESISTANCE IN INBRED

ECB1 = 3.75

ECB2 = 5.31530

Ear rate = 3.3125

#8 The comparable inbred to G3001 is ZS0560, an inbred having a number of similarities. ZS0560 is an inbred which has been or is presently in a number of commercial hybrids that are in a similar region of adaptation as most of the hybrids formed with G3001.

The Munsell code is a reference book of color, which is known and used in the industry and by persons with ordinary skill in the art of plant breeding.

The purity and homozygosity of inbred G3001 is constantly being tracked using isozyme genotypes as shown in Table 2.

## Isozyme Genotypes for G3001

Isozyme data were generated for inbred corn line G3001 according to procedures known and published in the art. The data in Table 2 gives the electrophoresis data on G3001 as compared to its two parents.

TABLE 2  
ELECTROPHORESIS RESULTS FOR G3001

	INBRED	ACP1	ACP4	ADH	MDH1	MDH2	PGD1	PGD2	PHI	PGM	IDH
5	G3001	33	55	22	22	11	22	22	22	22	22

#### Inbred and Hybrid Performance of G3001

10           The traits and characteristics of inbred corn line G3001 are listed to compare with other inbreds and/or in hybrid combination. The G3001 data shows the characteristics and traits of importance, giving a snapshot of G3001 in these specific environments.

15           Table 3A shows a comparison between G3001 and a comparable inbred ZS0560. G3001 has excellent vigor compared to inbred ZS0560. The two inbreds show significant differences in ear height, plant height, and across all Heat measurements for pollination and silking. G3001 has significantly higher yield at harvest than does ZS0560 but has slightly higher moisture. G3001 reaches heat  
20           peek with significantly less heat units than does ZS0560. The present invention is slightly less full season than is ZS0560. G3001 has significantly less medium round seeds and significantly more medium flat seeds than does ZS0560.

TABLE 3A G3001

YEAR	INBRED	VIGOR	EMERGE	PCT TILLER	PLANT HEIGHT	EAR HEIGHT	SHED	EAR QUALITY	PCT BARREN
OVERALL	G3001	6.4	84.9		174.7	63.3	3.7		
	ZS0560	5.5	84.0		165.2	68.9	4.3		
	# EXPTS	18	31		24	22	23		
	DIFF	0.9	0.8		9.5	5.6	0.6		
	PROB	0.001***	0.547		0.002***	0.016**	0.016		

YEAR	INBRED	HEATP10	HEATP50	HEATP90	HEATS10	HEATS50	HEATS90
OVERALL	G3001	1412	1456	1538	1407	1447	1476
	ZS0560	1464	1503	1606	1462	1502	1524
	# EXPTS	34	34	16	34	34	21
	DIFF	52	46	69	56	55	48
	PROB	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***

YEAR	INBRED	HEATPEAK	HEATBL	BL MOIST	% ROOT LODGE	% STALK LODGE	% DROP EARS	MOISTURE	YIELD
OVERALL	G3001	1350	2484					12.4	71.4
	ZS0560	1422	2578					11.6	66.5
	# EXPTS	27	2					37	37
	DIFF	71	94					0.5	4.8
	PROB	0.000***	0.010***					0.079*	0.023**

YEAR	INBRED	WARM GERM	COLD GERM	% LRG ROUND	% LRG FLAT	% MED ROUND	% MED FLAT	% SML ROUND	% SML FLAT
OVERALL	G3001	97.7	96.7			40.0	23.4		
	ZS0560	97.0	94.0			47.8	16.7		
	# EXPTS	21	21			11	11		
	DIFF	0.7	2.8			7.8	6.7		
	PROB	0.055*	0.002***			0.001***	0.001***		

\* .05 < PROB <= .10

\*\* .01 < PROB <= .05

\*\*\* .00 < PROB <= .01

\* .05 < PROB <= .10  
\*\* .01 < PROB <= .05  
\*\*\* .00 < PROB <= .01

YEAR	HYBRID	MATURITY	VM	FI
OVERALL	G3001 Hybrid		10.4	141
	CI Hybrid		9.3	138
	# EXPTS		145	142
	DIFF		1.0	2.6
	PROB		0.000***	0.002***

Table 3 b shows the inbred G3001 in hybrid combination with a common inbred and another hybrid combination that includes the same common inbred and another inbred. When in these hybrid combinations the present inbred carries less strength in field agronomic traits like lodging and ear drop then the comparison hybrid. However the G3001 hybrid combination brings distinctively significant reduction in moisture levels into the hybrid. The Y/M for the hybrid combination containing the present invention is significantly higher than the comparison hybrid.

Table 4 shows the GCA (general combining ability) estimates of G3001 compared with the GCA estimates of the other inbreds. The estimates show the general combining ability is weighted by the number of experiment/location combinations in which the specific hybrid combination occurs. The interpretation of the data for all traits is that a positive comparison is a practical advantage. A negative comparison is a practical disadvantage. The general combining ability of an inbred is clearly evidenced by the results of the general combining ability estimates. This data compares the inbred parent in a number of hybrid combinations to a group of "checks". The check data is from other companies' hybrids, particularly the leader in the industry and Garst Seed Company's commercial products and pre-commercial hybrids, which were grown in the same sets and locations.

TABLE 4

G3001

	N	FI	Y/M	GI	YLD	MST	%SL	%RL	%DE	TWT	POP	RM
XT=	26	1.1	0.5	0.1	3.8	0.4	-0.8	-1.1	-0.0	-1.2	-1	107

ZS0560

	N	FI	Y/M	GI	YLD	MST	%SL	%RL	%DE	TWT	POP	RM
XT =	60	0.2	-0.2	1.3	1.6	-0.5	0.5	0.0	0.0	-0.1	-209	109

$$FI = 100 + 0.5(Yld) - 2.3(MST) - 0.9(\%SL) - 0.9(\%RL) - 2.7(\%DE)$$

POP=plants per acre

RM=The Minnesota Relative Maturity

XT= GCA Estimate weighted by parent 2, but using only those parent 2 with two years of data

Table 4 shows G3001 in XT crossed to different inbreds to form 26 hybrid combinations. G3001 in hybrid combination shows an excellent advantage for moisture (MST) and an excellent advantage for yield compared to the commercial checks and the company's commercial inbreds. G3001 has a positive rating for yieldover moisture. In a number of categories the present invention surpasses the ZS0560 line. The ZS0560 line yield advantage is half of the present invention.

TABLE 5A

## YIELD RESPONSE

Research Plots

<u>HYBRID</u>	<u>YIELD</u>					
G3001/ Inbred X	66	94	123	152	180	209
Environment	75	100	125	150	175	200

Error: 12.7

#Plots 180

TABLE 5B  
YIELD RESPONSE

Research Plots

5	HYBRID	YIELD					
		71	96	120	145	169	194
	InbredX/inbred	71	96	120	145	169	194
	Environment	75	100	125	150	175	200

Error: 14.1

#Plots 2700

10

Table 5A shows the yield response of G3001 in hybrid combination in comparison with the plants in the environment around it at the same location. The data for the present inbred is showing consistently better results than the data of the comparison hybrid except in the lowest yielding environments. G3001 in hybrid combination, works well by providing yields better than the environment in high yielding environments. The yield is about equal to yields in the medium yielding environments. Its performance shows that this inbred is best suited to good environments.

15

20

Table 5b shows the data from a different hybrid that was formed with the same inbred X. This comparison hybrid is no longer yielding well particularly in the low and mid environments.

25

The present inbred also carries nice package of average to above average resistance levels to Eyespot, northern leaf blight and other diseases. This resistance allows the yield potential to be met more frequently and in more environments than many other inbreds.

The foregoing is set forth by way of example and is not intended to limit the

scope of the invention.

This invention also is directed to methods for producing a corn plant by crossing a first parent corn plant with a second parent corn plant wherein the first or second parent corn plant is an inbred corn plant from the line G3001. Further, both first and second parent corn plants can come from the inbred corn line G3001. A variety of breeding methods can be selected depending on the mode of reproduction, the trait, the condition of the germplasm. Thus, any such methods using the inbred corn line G3001 are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, haploid by such old and known methods of using stock material that induces haploids and anther culturing and the like. Additionally, this maize can, within the scope of the invention, contain: a mutant gene such as but not limited to sugary 1 or shrunken 1 or waxy or AE or imazethapyr tolerant (IT or IR <sup>TM</sup>) mutant gene; or transgenic genes such as but not limited to insect resistant genes such as *Bacillus thuringiensis* (Cry genes), or herbicide resistant genes such as Pat gene or Bar gene, EPSP, or disease resistant genes such as the Mosaic virus resistant gene, etc., or trait altering genes such as flowering genes, oil modifying genes, senescence genes and the like.

Various culturing techniques known to those skilled in the art, such as haploid, (stock six is a method that has been in use for twenty years and is well known to those with skill in the art), transformation, and a host of other conventional and unconventional methods are within the scope of the invention. All plants and plant cells produced using the inbred corn line are within the scope of this invention. The term transgenic plant refers to plants having genetic sequences, which are introduced into the genome of a plant by a transformation method and the progeny thereof.

Transformation Methods - are means for integrating new genetic coding

sequences into the plant's genome by the incorporation of these sequences into a plant through man's assistance.

5        Though there are a large number of known methods to transform plants, certain types of plants are more amenable to transformation than are others. Tobacco is a readily transformable plant. The basic steps of transforming plants including monocots are known in the art. These steps are concisely outlined in U.S. patent number 5,484,956 "Fertile Transgenic *Zea mays* Plants Comprising Heterologous DNA Encoding *Bacillus Thuringiensis* Endotoxin" issued January 16, 10        1996 and U.S. patent number 5,489,520 "Process of Producing Fertile *Zea mays* Plants and Progeny Comprising a Gene Encoding Phosphinothricin Acetyl Transferase" issued February 6, 1996.

15        Plant cells such as maize can be transformed by a number of different techniques. Some of these techniques which have been reported on and are known in the art include maize pollen transformation (See University of Toledo 1993 U.S. Patent No. 5,177,010); Biolistic gun technology (See U.S. patent number 5,484,956); Whiskers technology (See U.S. patent numbers 5,464,765 and 5,302,523); Electroporation; PEG on Maize; *Agrobacterium* (See 1996 article on transformation of maize cells in Nature Biotechnology, Volume 14, June 1996) 20        along with numerous other methods which may have slightly lower efficiency rates than those listed. Some of these methods require specific types of cells and other methods can be practiced on any number of cell types.

25        The use of pollen, cotyledons, meristems and ovum as the target issue can eliminate the need for extensive tissue culture work. However, the present state of the technology does not provide very efficient use of this material.

      Generally, cells derived from meristematic tissue are useful. Zygotic embryos can also be used. Additionally, the method of transformation of

meristematic cells of cereal is also taught in the PCT application WO96/04392. Any of the various cell lines, tissues, plants and plant parts can and have been transformed by those having knowledge in the art. Methods of preparing callus from various plants are well known in the art and specific methods are detailed in patents and references used by those skilled in the art. Cultures can be initiated from most of the above-identified tissue. The only true requirement of the transforming material is that it can form a transformed plant. The transgenic gene can come from various non-plant genes (such as; bacteria, yeast, animals, and viruses) along with being from animal or plants.

The DNA used for transformation of these plants clearly may be circular, linear, and double or single stranded. Usually, the DNA is in the form of a plasmid. The plasmid usually contains regulatory and/or targeting sequences which assists the expression of the gene in the plant. The methods of forming plasmids for transformation are known in the art. Plasmid components can include such items as: leader sequences, transit polypeptides, promoters, terminators, genes, introns, marker genes, etc. The structures of the gene orientations can be sense, antisense, partial antisense, or partial sense: multiple gene copies can be used.

The regulatory promoters employed can be constitutive such as CaMv35S (usually for dicots) and polyubiquitin for monocots or tissue specific promoters such as CAB promoters, etc. The prior art includes but is not limited to octopine synthase, nopaline synthase, CaMv19S, mannopine synthase promoters. These regulatory sequences can be combined with introns, terminators, enhancers, leader sequences and the like in the material used for transformation.

The isolated DNA is then transformed into the plant. Many dicots can easily be transformed with *Agrobacterium*. Some monocots are more difficult to transform. As previously noted, there are a number of useful transformation

processes. The improvements in transformation technology are beginning to eliminate the need to regenerate plants from cells. Since 1986, the transformation of pollen has been published and recently the transformation of plant meristems has been published. The transformation of ovum, pollen, and seedlings meristem greatly reduce the difficulties associated with cell regeneration of different plants or genotypes within a plant can present. Duncan, from at least 1985-1988 produced literature on plant regeneration from callus. Both inbred and hybrid callus have resulted in regenerated plants. Somatic embryogenesis has been performed on various maize tissues, which was once considered unusable for this purpose. The prior art clearly teaches the regeneration of plants from various maize tissues.

The most common method of transformation is referred to as gunning or microprojectile bombardment. This Biolistic process has small gold-coated particles coated with DNA shot into the transformable material. Techniques for gunning DNA into cells, tissue, callus, embryos, and the like are well known in the prior art.

After the transformation of the plant material is complete, the next step is identifying the cells or material, which has been transformed. In some cases, a screenable marker is employed such as the beta-glucuronidase gene of the *uidA* locus of *E. coli*. Then, the transformed cells expressing the colored protein are selected for either regeneration or further use. In many cases, a selectable marker identifies the transformed material. The putatively transformed material is exposed to a toxic agent at varying concentrations. The cells not transformed with the selectable marker, which provides resistance to this toxic agent, die. Cells or tissues containing the resistant selectable marker generally proliferate. It has been noted that although selectable markers protect the cells from some of the toxic affects of the herbicide or antibiotic, the cells may still be slightly effected by the toxic agent by having slower growth rates. If the transformed material was cell

lines then these lines are regenerated into plants. The cells' lines are treated to induce tissue differentiation. Methods of regeneration of cellular maize material are well known in the art since early 1982. European Patent Application, publication 160,390, describes tissue culture of corn, which can be used by those skilled in the art. The plants from the transformation process or the plants resulting from a cross using a transformed line or the progeny of such plants are transgenic plants.

Various techniques known to those skilled in the art, such as haploid, transformation, and a host of other conventional and unconventional methods are within the scope of the invention. All plants and plant cells produced using inbred corn line G3001 are within the scope of this invention. The invention encompasses the inbred corn line used in crosses with other, different, corn inbreds to produce (F1) corn hybrid seeds and hybrid plants. This invention includes cells, which upon growth and differentiation produce corn plants having the physiological and morphological characteristics of the inbred line G3001.

A deposit of at least 2500 seeds of this invention will be maintained by Garst Seed Company, 2369 330th Street, Slater, Iowa 50244. Access to this deposit will be available during the pendency of this application to the Commissioner of Patents and Trademarks and persons determined by the Commissioner to be entitled thereto upon request. All restrictions on availability to the public of such material will be removed upon issuance of a granted patent of this application by depositing at least 2500 seeds of this invention at the American Type Culture Collection. The deposit of at least 2500 seeds will be from the same inbred seed taken from the deposit maintained by Garst Seed Company. The ATCC deposit will be maintained in that depository, which is a public depository, for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced if it becomes nonviable during

that period.

Additional public information on some ZS designations may be available from the PVP office a division of the US government.

5

Accordingly, the present invention has been described with some degree of particularity directed to the preferred embodiment of the present invention. It should be appreciated, though, that the present invention is defined by the following claims construed in light of the prior art so that modifications or changes may be made to the preferred embodiment of the present invention without departing from the inventive concepts contained herein.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206